arXiv:nlin/0612007v1 [nlin.CD] 4 Dec 2006

Conjugation of cascades

Jesús San Martín[a,b], Daniel Rodríguez-Pérez[b]

February 5, 2008

[a] Departamento de Matemática Aplicada, E.U.I.T.I, Universidad Politécnica de Madrid. Ronda de Valencia 3, 28012 Madrid Spain

[b] Departamento de Física Matemática y Fluidos, U.N.E.D. Senda del Rey 9, 28040 Madrid Spain

Corresponding author: jsm@dfmf.uned.es

Abstract

Presented in this work are some results relative to sequences found in the logistic equation bifurcation diagram, which is the unimodal quadratic map prototype. All of the different saddle-node bifurcation cascades, associated to every last appearance p-periodic orbit ($p = 3, 4, 5, \ldots$), can also be generated from the very Feigenbaum cascade. In this way it is evidenced the relationship between both cascades. The orbits of every saddle-node bifurcation cascade, mentioned above, are located in different chaotic bands, and this determines a sequence of orbits converging to every band-merging Misiurewicz point. In turn, these accumulation points form a sequence whose accumulation point is the Myrberg-Feigenbaum point. It is also proven that the first appearance orbits in the n-chaotic band converge to the same point as the last appearance orbits of the $(n+1)$-chaotic band. The symbolic sequences of band-merging Misiurewicz points are computed for any window.

1 Introduction

Nonlinear dynamical systems exhibit a rich variety of behaviors. Bifurcations, chaos, patterns, phase transitions are among the most investigated phenomena, and they have led to many discussions within scientific literature. The former mentioned phenomena appears in a lot of dynamical systems, and this is an indicative of the existence of underlaying common structures whose origin claims to be uncovered.

To be able to undertake the study of a dynamical system, diverse tools have been developed, which make the study of such problems easier, by means of simplifications, classifications or through the identification of universality classes among transitions from one behavior to another.

Near a hyperbolic fixed point, the well known Hartman-Großman theorem states that the local phase portrait is topologically conjugate to the phase portrait of the linearized system. However, when the fixed point is a non-hyperbolic one, the center manifold must be considered. The Center Manifold theorem allows one to generalize the ideas of the Hartman-Großman theorem. It is deduced that the flow, restricted to the center manifold, determines the topological behavior close to a non-hyperbolic fixed point. In general, when the flow is restricted to the center manifold the problem is simplified quite a bit, yet the restricted flow itself can be very complicated.

The normal form theorem is the next mathematical tool that must be used: a smooth vector field is substituted by a polynomial vector field. Another technique very similar, in certain way, to the tool stated above consists in constructing a Poincaré section. After that, the Poincaré map is created, that is, a map from the Poincaré section in itself. In dissipative systems this map contracts the Poincaré section surface after each iteration, and the initial problem is thus transformed into another problem with an attractor of lower dimensionality.

Fortunately, one-dimensional maps provide a good approximation to many dynamical systems. Furthermore one-dimensional maps can be classified into universality classes. That is why the bifurcation route to chaos, discovered by Feigenbaum [1, 2] and characterized by its universal scaling is so important. And for the same reason the logistic equation

$$x_{n+1} = f(x_n) = \mu x_n(1 - x_n) \quad \mu \in [0,4],\ x_n \in [0,1] \tag{1}$$

is so intensively studied. The different behaviors of this map can be extended to any quadratic one-dimensional map, because they are topologically conjugated [3]. Therefore, any dynamical system ruled by a quadratic polynomial behaves as if it were ruled by the logistic equation.

The goal of this paper is to show in the logistic equation the interesting connection between the Feigenbaum cascade and the saddle-node cascade [4]. Also this connection affects the global structures of windows and Misiurewicz points [5] where chaotic bands mix. Given the multitude of dynamical systems that exhibit Feigenbaum's universality, it is very probable that they also show the new universal behaviors described in this paper.

Periodic windows, apart from being an irreplaceable tool in system dynamics, also play other roles and appear in different problems of nonlinear dynamics. They can be used, by applying the Poincaré map, to spot the windows of a dynamical systems which are ruled by ordinary differential equations [6]. Chaos control techniques use periodic windows too and they use unstable periodic orbits inside them [7, 8]. Periodic windows are also useful to study chaos-induced diffusion [9] which is affected by the presence of proximate periodic windows. Therefore any contribution aimed at the identification of the laws ruling periodic windows will be useful in these related fields.

This paper is organized as follows:

In section 3, it is proven that there is an analytic relation among saddle-node bifurcation cascades [4] and Feigenbaum cascade [1, 2]. The former is derived from the latter. We will refer to this relation by *conjugation of cascades*. The section 3 is preceded by section 2 where later used definitions and theorems are introduced.

In the following section 4, the previous result is used to prove that saddle-node bifurcation cascades associated to last appearance orbits in the $(n+1)$-chaotic band and the ones associated to first appearance orbits in the n-chaotic band converge to the same Misiurewicz point where the $(n+1)$-chaotic and n-chaotic bands mix.

Section 5 generalizes the former results, which were only valid in the canonical window, to any periodic window. In particular, we will also generalize the rule to get the saddle-node bifurcation cascade in any window. As a spin-off, a geometrical interpretation of the Derrida, Gervois and Pomeau (DGP) formula [10] is achieved.

In Section 6, some potential applications of the theorems to experimental research are indicated.

2 Definitions

The following notation, definitions and theorems will be referred to in what follows. Whenever they are based on or copied from a previous work, the original reference is given.

Definition 1. S_n will denote a n-periodic symbolic pattern. The symbols are R or L, depending on wether the iterates x_i of a n-periodic orbit of the logistic map (1) are $x_i > \frac{1}{2}$ or $x_i < \frac{1}{2}$ $(i = 1, 2, 3, \dots n)$, respectively [11]. In particular, to determine in an unique way the periodic sequence S_n, we will always take as the first letter in the sequence that corresponding to the position x_i nearest to $1/2$, the rest of the symbols being determined by the succesive iterations of this first value [4].

Definition 2. The pattern S_n has even "R-parity" if it has an even number of symbols R in it. Otherwise S_n "R-parity" will be termed odd [11]. We will use R-parity whenever we work with logistic-like maps (upwards convex). Should we were working with the real Mandelbrot map, for instance (which is downwards convex), all R's and L's in what follows would appear interchanged, and we would use L-parity instead.

Definition 3. The pattern $\overline{S_n}$ obtained exchanging R and L in S_n will be called the conjugated sequence of S_n. In the case of a supercycle pattern CS_{n-1} (C corresponds to $x_1 = 1/2$), its conjugate will only change the R and L in the pattern, whereas the initial C remains in place.

Definition 4. $(S_n)^p = \underbrace{S_n S_n \ldots S_n}_{p \text{ times}}$, is the pattern of $n \cdot p$ symbols constructed by p repetitions of the n-symbols pattern S_n. As a consequence, we may write $[(S_n)^p]^q = (S_n)^{p \cdot q}$. We will adopt an analogous notation for the case of supercycles.

Definition 5. CS_{n-1} is a n-symbols pattern formed by replacing the first letter in sequence S_n by the symbol C. This orbit corresponds to a supercycle periodic orbit (starting at $x_1 = \frac{1}{2}$).

Definition 6. $(CS_n)^p|S_p$ is the pattern resulting from the substitution of the p succesive C-letters in the pattern $(CS_n)^p = \underbrace{CS_n CS_n \ldots CS_n}_{p \text{ times}}$ by the p succesive symbols in S_p.

Definition 7. $CP_{n,q}$ represents the pattern of the n-th pitchfork bifurcation supercycle of the q-periodic orbit. For $q = 1$ we will write simply CP_n.

Definition 8. The first F-harmonic of CS_n, denoted by $H_F^{(1)}(CS_n)$, is formed appending CS_n to itself and changing the second C to R (L) if the R-parity of CS_n is even (odd). The second F-harmonic, $H_F^{(2)}(CS_n)$, is formed appending CS_n to $H_F^{(1)}(CS_n)$ and changing the new C to R (L) is the R-parity of $H_F^{(1)}(CS_n)$ is even (odd). The successive F-harmonics are constructed in the same way. [12]

The F-harmonics (or Fourier harmonics) were introduced by Romera, Pastor and Montoya in [12] and have been widely employed by these authors in the study of dynamical systems [13, 14, 15]. It must be emphasized that, while those harmonics introduced by Metropolis, Stein and Stein [11] give us the patterns corresponding to the Feigenbaum period doubling cascade, the F-harmonics are used to compute the patterns of the last appearance orbits (see Definition 10 below) in the current window. This property will be amply used in our proofs; they will be, together with the saddle-node bifurcation cascades, the main tools used in this paper.

Definition 9. A saddle-node bifurcation cascade is a sequence of saddle-node bifurcations in which the number of fixed points showing this kind of bifurcation is duplicated [4]. The successive elements of the sequence are given by an equation identical to the one that Feigenbaum found for a period doubling cascade [1, 2]. This implies that both bifurcation cascades (Feigenbaum's and saddle-node) scale in the same way.

Definition 10. We will consider a q-periodic window to be a first appearance window within the p-periodic one if it corresponds to values of parameter μ of expression (1) smaller than any other q-periodic window inside the given p-periodic one. Similarly, we will define a last appearance q-periodic orbit (within the p-periodic one) as that having the larger μ parameter among the period-q windows (of the p-periodic one).

A periodic window of period $q \cdot p$ will be called the q-periodic window inside the given p-periodic one, to stress the similarity of the structures in period-p window with those in the canonical window.

Definition 11. Within the chaotic region of the logistic equation bifurcation diagram (to the right of the Myrberg-Feigenbaum point), there exist chaotic bands of μ values where the iterates of expression (1) tend to be grouped in 2^n intervals (separated by $2^n - 1$ "empty" intervals); we will refer to one of these bands as the n-th chaotic band. Following the same naming convention as in Definition 10, we will speak of the first and last appearance $q \cdot 2^n$-periodic orbits inside a chaotic band, as those corresponding to the smaller and greater parameter μ values, respectively, within that n-chaotic band.

In our proofs we will often use saddle-node cascades inside the canonical window. They can be worked out applying the following theorem, which is a particular case of that shown in [4].

Theorem 1 (*Saddle-Node bifurcation cascade in the canonical window*). *The sequence of the $p \cdot 2^n$-periodic saddle-node orbit of a saddle-node bifurcation cascade starting from a p-periodic orbit in the canonical window is obtained with the following process:*

(i) *Write the sequence of the orbit of the supercycle of f^{2^n}, that is CP_n (see Definition 7).*

(ii) *Write consecutively p times the sequence obtained in the former point "i", getting a sequence like*

$$\underbrace{CP_n \dots CP_n}_{p \text{ times}}$$

(iii) *Write the sequence of period-p saddle-node orbit, that is, the sequence of the saddle-node orbit of f^p. The first point in the sequence must be the nearest saddle-node point to C.*

(iv) *If n is odd then conjugate the letters obtained in point "iii" by means of $L \leftrightarrow R$. Bear in mind that $n \in \mathcal{Z}^+$.*

(v) *Replace the i-th letter C ($i = 1, 2, \dots p$) of the sequence obtained in "ii" by the i-th letter of the sequence obtained in "iv".*

3 Conjugation of cascades.

Theorem 2. (*Conjugation of cascades in the canonical window*) *Let $S_{p \cdot 2^n}$ be the pattern of the $p \cdot 2^n$-periodic orbit of a saddle-node bifurcation cascade in the canonical window. If $CS_{p \cdot 2^n - 1}$ denotes the supercycle associated to $S_{p \cdot 2^n}$ (see Definition 5) and CP_n denotes the supercycle of the n-th pitchfork bifurcation (see Definition 7) then*

$$CS_{p \cdot 2^n - 1} = H_F^{(p-1)}(CP_n)$$

That is, the supercycle of the n-th element of the cascade can be computed from the saddle-node bifurcation cascade (using Theorem 1) or from the Feigenbaum cascade (using F-harmonics, see Definition 8).

Proof. To prove this statement, we are going to compute the pattern of one of these $2^n p$-periodic orbits in two ways: 1) applying the **Theorem 1** *(Saddle-Node bifurcation cascade in the canonical window),* this way we get the n-th term of the saddle-node bifurcation cascade, and 2) computing the $(p-1)$-th F-harmonic to the 2^n-periodic orbit generated after the n-th pitchfork bifurcation of the 1-periodic orbit.

In our proof we must consider separately the cases of even and odd n.

Let n be even:

(i) Applying Theorem 1:

Theorem 1 involves the following steps:

- determine the last apperarance p-periodic orbit pattern, that is CRL^{p-2} [16]. Given that n is even, this pattern will not be conjugated (i.e. R's and L's will not be swapped).
- repeat p times the sequence CP_n which gives
$$\underbrace{CP_nCP_n\ldots CP_n}_{p\text{ times}} \tag{2}$$
- substitute the succesive C symbols in the pattern (2) by the succesive symbols in the pattern CRL^{p-2}, which gives the pattern sought for:
$$CP_nRP_n(LP_n)^{p-2} \tag{3}$$
where $p = 3,\ 4,\ 5, \ldots$ (there is no cascade for $p = 2$).

(ii) Applying F-harmonics:

The construction of the F-harmonics follows the next steps:

- append CP_n to the pattern CP_n, giving
$$CP_nCP_n$$
- substitute the second C by R (or L) if the R-parity of CP_n is even (or odd). Given that n is assumed to be even, the R-parity of CP_n is also even (see Note 1 in Appendix); as a consequence the first F-harmonic becomes
$$H_F^{(1)}(CP_n) = CP_nRP_n$$
- generate the n-th F-harmonic appending CP_n to the $(n-1)$-th F-harmonic, and changing the second C of the obtained sequence by R (or L) if the R-parity of $H_F^{(n-1)}(CP_n)$ is even (or odd). It is easy to see that $H_F^{(1)}(CP_n)$ R-parity is odd. As a consequence, on

appending the expression CP_n to $H_F^{(1)}(CP_n)$ in order to obtain the $H_F^{(2)}(CP_n)$, the C will be changed to L, thus conserving R-parity. The same will happen for the next F-harmonics thus getting

$$H_F^{(p-1)}(CP_n) = CP_nRP_n(LP_n)^{p-2} \tag{4}$$

where $p = 2,\ 3,\ 4,\ldots$ (although in the cascade there is no $p = 2$).

Let n be odd:

(iii) Applying Theorem 1:

In this case of n odd, Theorem 1 would be equally applied but pattern CRL^{p-2} must be conjugated to CLR^{p-2}. Thus the final pattern will be

$$CP_nLP_n(RP_n)^{p-2} \tag{5}$$

(iv) Applying F-harmonics:

Being n odd, the R-parity of CP_n is also odd (see Note 1 in Appendix), thus modifying the first F-harmonic to

$$H_F^{(1)}(CP_n) = CP_nLP_n$$

having even R-parity. As a consequence, on appending CP_n to $H_F^{(1)}(CP_n)$ to get to $H_F^{(2)}(CP_n)$ the second C will be replaced by R thus conserving the R-parity. The result is

$$H_F^{(p-1)}(CP_n) = CP_nLP_n(RP_n)^{p-2}$$

with $p = 2,\ 3,\ 4,\ 5,\ldots$

In summary, the theorem is proven because both Theorem 1 and the F-harmonics give the same results

$$\begin{array}{ll} CP_nRP_n(LP_n)^{p-2} & p = 3,4,5,\ldots \text{ for } n \text{ even} \\ CP_nLP_n(RP_n)^{p-2} & p = 3,4,5,\ldots \text{ for } n \text{ odd} \end{array} \tag{6}$$

The terms with $p = 2$ in the F-harmonic have been disregarded, given that they correspond to a pitchfork bifurcation which, thus, cannot account for a saddle-node one.◀

Examples.

As an example, we will show how to obtain the first saddle-node bifurcation orbits of the last appearance period-5 orbit in the canonical window, whose pattern is CRL^3. It will be enough to examine the cases $n = 1, 2$ to show how to perform the calculation in both cases when the P_n orbit has to be conjugated or not.

For $n = 1$, that is odd n:

- Using Theorem 1 we compute:
    - the corresponding CP_1 in the canonical window is simply CR
    - the pattern CRL^3 will be conjugated because n is odd, resulting: $CRL^3 \to CLR^3$
    - repeat CR $p = 5$ times: $CRCRCRCRCR$
    - replace C's in the latter sequence by CLR^3, resulting: $CRLR^7$
- Using Theorem 2 and F-harmonics:
    - $CP_1 = CR$
    - its first F-harmonic is $CRCR \to CRLR$ (C changes to L because R-parity of CR is odd)
    - its second F-harmonic is $CRLRCR \to CRLR^3$ (C changes to R because R-parity of $CRLR$ is even)
    - and its fourth F-harmonic is $CRLR^7$

Both sequences coincide, as it was expected.

For $n = 2$, that is even n:

- Using Theorem 1 we compute:
    - $CP_2 = CRLR$
    - the pattern CRL^3 will not be conjugated because n is even
    - repeat $CRLR$ $p = 5$ times: $CRLRCRLRCRLRCRLRCRLR$
    - replace C's in the latter sequence by CRL^3, resulting: $CRLR^3(LR)^7$
- Using Theorem 2 and F-harmonics:
    - $CP_2 = CRLR$

  - its first F-harmonic is $CRLRCRLR \to CRLR^3LR$ (C changes to R because R-parity of $CRLR$ is even)
  - and its fourth F-harmonic is $CRLR^3(LR)^7$

Both sequences also coincide.

4 Convergence of first and last appearance orbits.

Theorem 3. *In the canonical window, the last appearance orbits within the $(n+1)$-chaotic band have the same accumulation point as the first appearance orbits within the n-chaotic band.*

Proof. To prove the theorem, we need to compute the patterns of the first and last appearance orbits within an arbitrary chaotic band. These patterns are given by the saddle-node bifurcation cascades associated, respectively, to the first and last orbits in the 1-chaotic band. The saddle-node cascades will be computed by means of Theorem 1. Given that this theorem treats the cases of even and odd n differently, we will study these two cases separately.

Let n be even:

(i) Computation of the first appearance orbit patterns within the n-chaotic band.

As we have already indicated, we compute the requested pattern applying Theorem 1 following the next steps:

- give the pattern of the first appearance superstable p-periodic orbit within the 1-chaotic band. This pattern is $CRLR^{p-3}$, with $p = 3, 5, 7, \ldots$
- repeat p times the pattern CP_n (see Definition 7) as in the following

$$\underbrace{CP_nCP_n \ldots CP_n}_{p \text{ times}} \tag{7}$$

Given that we assume n to be even, the symbols in the pattern $CRLR^{p-3}$ are not conjugated, and succesive C's in expression (7) are simply replaced by the successive C's in the pattern $CRLR^{p-3}$. This substitution results in the sought for pattern

$$CP_nRP_nLP_n(RP_n)^{p-3} \tag{8}$$

where $p = 3, 5, 7, \ldots$ As expected, the pattern (8) has $p \cdot 2^n$ symbols.

(ii) Computation of the last appearance orbit patterns within the $(n+1)$-chaotic band.

To apply Theorem 1 we accomplish the following steps:

- give the pattern of the last appearance superstable p-periodic orbit within the 1-chaotic band, that is [16]

$$CRL^{p-2}$$

with $p = 3,\, 4,\, 5, \ldots$

- repeat p times the pattern CP_{n+1}. Given that n is even, $CP_{n+1} = CP_nRP_n$ (see Note 1 in Appendix), this becomes

$$\underbrace{CP_nRP_nCP_nRP_n \ldots CP_nRP_n}_{p \text{ times}} \tag{9}$$

- given that $n+1$ is odd the pattern CRL^{p-2} will be conjugated, resulting in CLR^{p-2}. Next, succesive C's in expression (9) are replaced by successive symbols in CLR^{p-2}. This substitution gives the sought for pattern, that is

$$CP_nRP_nLP_n(RP_n)^{2p-3}$$

where $p = 3,\, 4,\, 5, \ldots$

Let n be odd:

(iii) Computation of the first appearance orbit patterns within the n-chaotic orbit.

We take an identical approach to "i", but given that n is odd, the pattern $CRLR^{p-3}$ must be conjugated to $CLRL^{p-3}$. The consequence of this conjugation is that the following pattern is generated

$$CP_nLP_nRP_n(LP_n)^{p-3}$$

with $p = 3,\, 5,\, 7, \ldots$

(iv) Computation of the last appearance orbit patterns within the $(n+1)$-chaotic orbit.

We take an identical approach to "ii", but given that $(n+1)$ is even, the pattern CRL^{p-2} must not be conjugated. On the other hand, given that n is odd, we have $CP_{n+1} = CP_nLP_n$ (see Note 1 in Appendix). As a consequence, the pattern

$$CP_nLP_nRP_n(LP_n)^{2p-3}$$

is obtained, with $p = 3,\ 4,\ 5, \ldots$

In summary we get:

- the first appearance orbit patterns within the n-chaotic band are:

$$\begin{array}{lll} CP_nRP_nLP_n(RP_n)^{p-3} & p = 3,5,7,\ldots & \text{for } n \text{ even} \\ CP_nLP_nRP_n(LP_n)^{p-3} & p = 3,5,7,\ldots & \text{for } n \text{ odd} \end{array} \tag{10}$$

- the last appearance orbit patterns within the $(n+1)$-chaotic band are:

$$\begin{array}{lll} CP_nRP_nLP_n(RP_n)^{2p-3} & p = 3,4,5,\ldots & \text{for } n \text{ even} \\ CP_nLP_nRP_n(LP_n)^{2p-3} & p = 3,4,5,\ldots & \text{for } n \text{ odd} \end{array} \tag{11}$$

Myrberg's formula [17] shows that the parameters corresponding to these orbits have the same accumulation point, and this proves the theorem. ◀

4.1 Consequences of Theorem 3 (Misiurewicz point cascade)

The limit $p \to \infty$ of expressions (10) and (11) corresponds to those points where the n and $n+1$ bands merge. These are Misiurewicz points having a preperiod (within square brackets in the following expression) followed by a period that corresponds to an unstable periodic orbit:

$$\begin{array}{ll} [CP_nRP_nLP_n]\ RP_n & \text{for } n \text{ even} \\ [CP_nLP_nRP_n]\ LP_n & \text{for } n \text{ odd} \end{array} \tag{12}$$

These expressions were previously obtained in [13] using a different approach. The value of the parameter μ in expression (1), where these points are localized, is the smallest solution of the equation

$$\mathcal{P}_{2^n+1}(\mu) = \mathcal{P}_{2^{n-1}}(\mu)$$

being $\mathcal{P}_n(\mu)$ the dynamical polynomials defined in [17, 18], which can be obtained recursively (for equation (1)) as

$$\mathcal{P}_0 = 1/2, \quad \mathcal{P}_{n+1} = \mu\mathcal{P}_n(1 - \mathcal{P}_n)$$



Figure 1: First and last appearance orbits approaching the band merging Misiurewicz point.

Given that patterns (12) have been obtained as the limit of saddle-node orbit sequences elements (each of these elements belongs to a different saddle-node bifurcation cascade) the following corollary can be stated:

Corollary to Theorem 3. *Band merging points, whose patterns are given in (12) as the limit orbits of saddle-node orbit sequences, form a Misiurewicz point cascade. Furthermore, this Misiurewicz point cascade converges to the Myrberg-Feigenbaum point, driven by the behavior of the saddle-node bifurcation cascades, as is described in [4].*

5 Generalization to any q-periodic window

The selfsimilarity shown by the behaviors of the logistic equation is well known. It is revealed by the existence of periodic windows that mimic the different structures of the canonic window. Inside every subwindow the same bifurcation diagram structure exists, with its periodic and chaotic regions, intermittencies, and so on. In particular, every window shows in its beginning a periodic behavior, followed by a pitchfork bifurcation cascade; similarly every window has saddle-node bifurcation cascades, which reflect the ones existing in the canonical window.

This circumstance is a clear suggestion to look for the relationship between the orbits of the pitchfork bifurcation cascade and those of the saddle-node bifurcation cascade, both belonging to a given q-periodic window. In this way we generalize the results obtained above, for the canonical window, to any arbitrary window. Nonetheless, although the results for the canonical window will become a particular case of the more general theorems, they will be used in the proofs of the general theorems.

The Bifurcation Rigidity theorem [19] allows us to approximately compute the parameter values for which the different orbits in a p-periodic window, mimicking those in the canonical one, take place. This is achieved through a linear mapping of the windows, the deeper the subwindow is located the more exact the result is. Nevertheless, this useful tool does not allow us to determine the patterns of the orbits and thus, we cannot establish a relationship among them.

To obtain the patterns of the orbits, we need to compute that of the $p \cdot 2^n$-periodic orbits in a saddle-node bifurcation cascade, located inside a q-periodic window. A similar problem (that of finding the pattern of a q-periodic orbit inside the $p \cdot 2^n$-periodic window) was already addressed in [4] and is thus known to us. Merely exchanging the names of the windows involved, brings us the requested pattern. This result, using the compact notation introduced in section 2, is stated as follows.

Theorem 4. *Let $S_{p\cdot 2^n}$ ($n = 1, 2, 3\ldots$) be the patterns of the orbits in a saddle-node bifurcation cascade located inside the canonical window. Let S_q be the pattern of the q-periodic saddle-node orbit which marks the origin of the q-periodic window, and let CS_{q-1} be the pattern of its associated supercycle. The pattern $S_{q\cdot p\cdot 2^n}$ of the $p \cdot 2^n$-periodic orbit in the saddle-node bifurcation cascade inside the q-periodic window considered is*

$$\big((CS_{q-1})^{p\cdot 2^n}|S_{p\cdot 2^n}\big) \qquad \textit{for even R-parity of the q-periodic supercycle}$$

$$\big((CS_{q-1})^{p\cdot 2^n}|\overline{S_{p\cdot 2^n}}\big) \qquad \textit{for odd R-parity of the q-periodic supercycle}$$

Proof. The proof is achieved by just exchanging $f^q \leftrightarrow f^{p\cdot 2^n}$ in the original proof (see [4]). With this substitution, expression

$$\big((CS_{q-1})^{p\cdot 2^n}|S_{p\cdot 2^n}\big)$$

is obtained, which is valid for f^q having a maximum at $x = 1/2$. This corresponds to an even R-parity of the q-periodic orbit (see Lemma 1 in the appendix). Should f^q have a minimum at $x = 1/2$, that is, for a q-periodic orbit having odd R-parity (see Lemma 2 in the appendix), it is enough to conjugate $S_{p\cdot 2^n} \rightarrow \overline{S_{p\cdot 2^n}}$ to get the correct result, that is

$$\big((CS_{q-1})^{p\cdot 2^n}|\overline{S_{p\cdot 2^n}}\big)$$

◀

As a consequence of replacing $f^q \leftrightarrow f^{p\cdot 2^n}$, we observe that the orbit of $f^{p\cdot 2^n}$ gets reproduced around every extremum of f^q lying near the line $x_{i+1} = x_i$. This is shown in figure 2, where the shape of the f^3 orbit (upper panel in figure 2) can be identified in those extremes of f^5 (right panel in figure 2) near the diagonal $x_{i+1} = x_i$. In figure 3 the same situation is shown, this time for the supercycle of $f^{5\cdot 3\cdot 2}$ (saddle-node bifurcation of that of $f^{5\cdot 3}$); in this case it is $f^{3\cdot 2}$ that gets reproduced in each maximum of f^5.

Let us emphasize that, what is being obtained is the pattern of a $p \cdot 2^n$-periodic saddle-node orbit located inside a q-periodic window. Given that this q-periodic window has its origin at a q-periodic saddle-node orbit, the resulting pattern will be that of a $q \cdot p \cdot 2^n$-periodic saddle-node orbit. Moreover, the geometrical construction used in the proof, allows us to show the geometrical meaning of the DGP formula, as can be seen in what follows.



Figure 2: Graph of $f^{5\cdot3}$ (the f^3 periodic orbit inside the period-5 window). It can be seen how in one of the extrema of f^5 (highlighted in the right panel plot) the shape of f^3 is reproduced.



Figure 3: Graph of $f^{5\cdot3\cdot2}$ (the $f^{3\cdot2}$ saddle-node bifurcation of the f^3 periodic orbit inside the period-5 window, which is located in the 1-chaotic band of the period-5 window). It can be seen how in one of the extrema of f^5 (drawn in the right panel and encircled by the ellipse) the shape of $f^{3\cdot2}$ is reproduced.

The substitution of the first symbol in the pattern $S_{q\cdot p\cdot 2^n}$ by a C is enough to obtain its associated supercycle, that we denote by $CS_{q\cdot p\cdot 2^n-1}$ (see Definition 5), whose geometrical interpretation was given above. Given that this same supercycle can be also constructed applying DGP rule to compute the product of the orbits $CS_{p\cdot 2^n-1}$ and CS_{q-1}, what is obtained is a geometrical interpretation of the DGP rule. We thus conclude (see figure 4 for a graphical sketch) that:

1. The composition of the supercycles $CS_{p\cdot 2^n-1}$ and CS_{q-1} , according to DGP rule, represents the supercycle associated to the $p \cdot 2^n$-periodic orbit, inside the q-periodic window.

   That is so because, in the geometric proof, we have forced the shape of a $p \cdot 2^n$-periodic saddle-node orbit to appear around the extrema of a saddle-node q-periodic orbit. This is shown in figures 2 and 3.

2. However, we could have forced the reverse situation, that is, to reproduce the shape of the q-periodic orbit around the extrema of a $p\cdot 2^n$-periodic orbit, arising after n pitchfork bifurcations of a p-periodic saddle-node orbit. In that case, we would be focusing on the n-th pitchfork bifurcation of a p-periodic orbit, localized inside the q-periodic window. This second case was already known and is not a conclusion of our geometrical proof.

Theorem 4 allows us to state Theorem 2 and Theorem 3 generalized to subwindows inside the canonical one. The proofs are given in the next two subsections.

5.1 Conjugation of cascades in an arbitrary window.

Theorem 5 (*Conjugation of cascades in an arbitrary window*). *Let $S_{q\cdot p\cdot 2^n}$ be the pattern of the $p \cdot 2^n$-periodic orbit in the saddle-node bifurcation cascade inside the q-periodic window, such that p is a last appearance orbit. If $CS_{q\cdot p\cdot 2^n-1}$ denotes the supercycle associated to $S_{q\cdot p\cdot 2^n}$ (see Definition 5) and $CP_{n,q}$ denotes the n-th pitchfork bifurcation supercycle of the q-periodic orbit (see Definition 7) then*

$$CS_{q\cdot p\cdot 2^n-1} = H_F^{(p-1)}(CP_{n,q})$$

What the theorem says is that, inside an arbitrary q-periodic window, superstable orbit patterns of a saddle-node bifurcation cascade, which



Figure 4: Example of the application of Theorem 5, within the period-3 window to find the last appearance orbits in the 1- and 2-chaotic bands, applying F-harmonics to the orbits in the Feigenbaum cascade. Also shown, is how the Feigenbaum cascade is computed from Metropolis-Stein-Stein harmonics [11], and the saddle-node bifurcation cascade is obtained using Theorem 4.

is associated to a last appearance orbit, coincide with orbit patterns generated by appying F-harmonics to Feigenbaum cascade supercycles.

Proof. Let us consider a q-periodic window, whose origin is in a q-periodic saddle-node orbit. This q-periodic window needs not to be a primary window, any window located inside any other will do.

Let us also consider the last appearance p-periodic orbit inside the chosen q-periodic window, where $p = 3,\, 4,\, 5, \ldots$. This p-periodic orbit has a saddle-node bifurcation cascade associated to it, whose orbits have periods $p \cdot 2^n$, $n = 0,\, 1,\, 2, \ldots$, and are all located inside the q-periodic window (that is, they are really $q \cdot p \cdot 2^n$-periodic orbits).

We want to show that the $p \cdot 2^n$-periodic saddle-node orbit supercycle inside the q-periodic window can be obtained equivalently by applying Theorem 4 or computing the $(p-1)$-th F-harmonic of the n-th pitchfork bifurcation of the q-periodic orbit.

Let the R-parity of the q-periodic supercycle pattern be even.

(i) Applying Theorem 4.

This theorem (see section 5) says that the pattern $CS_{q \cdot p \cdot 2^n - 1}$ is given by

$$\left((CS_{q-1})^{p \cdot 2^n} | CS_{p \cdot 2^n - 1}\right) \text{ for even } R\text{-parity of the } q\text{-periodic orbit}$$

On the other hand, the pattern $CS_{p \cdot 2^n - 1}$ is given by expression (6).

(a) Let n be odd. Then $CS_{p \cdot 2^n - 1} = CP_n LP_n (RP_n)^{p-2}$ according to expression (6). Then we perform the following computation

$$\begin{aligned} \left((CS_{q-1})^{p \cdot 2^n} | CS_{p \cdot 2^n - 1}\right) &= \left((CS_{q-1})^{p \cdot 2^n} | CP_n LP_n (RP_n)^{p-2}\right) \\ &= \left((CS_{q-1})^{2^n} | CP_n\right) \left((CS_{q-1})^{2^n} | LP_n\right) \\ &\quad \left((CS_{q-1})^{2^n} | RP_n\right)^{p-2} \end{aligned}$$

(b) Let n be even. Then $CS_{p \cdot 2^n - 1} = CP_n RP_n (LP_n)^{p-2}$, with $p = 3,\, 4,\, 5, \ldots$ according to expression (6). In this case the computation goes as follows

$$\begin{aligned} \left((CS_{q-1})^{p \cdot 2^n} | CS_{p \cdot 2^n - 1}\right) &= \left((CS_{q-1})^{p \cdot 2^n} | CP_n RP_n (LP_n)^{p-2}\right) \\ &= \left((CS_{q-1})^{2^n} | CP_n\right) \left((CS_{q-1})^{2^n} | RP_n\right) \\ &\quad \left((CS_{q-1})^{2^n} | LP_n\right)^{p-2} \end{aligned}$$

(ii) Applying F-harmonics.

Let us now apply F-harmonics to the n-th pitchfork bifurcation of the q-periodic orbit.

The n-th pitchfork bifurcation of CS_{q-1} , that is $CP_{n,q}$, is given by

$$\left((CS_{q-1})^{2^n}|CP_n\right)$$

and has the same R-parity as CP_n, according to Lemma 2 in the Appendix.

(a) Let n be odd, then the R-parity of CP_n is odd and, hence, the R-parity of $\left((CS_{q-1})^{2^n}|CP_n\right)$ is also odd (for $n \geq 1$). The F-harmonic is computed as

$$\begin{aligned} &H_F^{(1)}\left[(CS_{q-1})^{2^n}|CP_n\right] = \\ &= \left[(CS_{q-1})^{2^n}|CP_n\right]\left[(CS_{q-1})^{2^n}|LP_n\right] \end{aligned}$$

where the C's in the last bracket have been changed to L's because of the assumed odd R-parity. As a consequence, $H_F^{(1)}\left[(CS_{q-1})^{2^n}|CP_n\right]$ has even R-parity. To compute the second F-harmonic $H_F^{(2)}\left[(CS_{q-1})^{2^n}|CP_n\right]$, those C's will be replaced by R thus keeping the same R-parity. Hence, the result for the $(p-1)$-th F-harmonic is

$$\begin{aligned} &H_F^{(p-1)}\left[(CS_{q-1})^{2^n}|CP_n\right] = \\ &= \left[(CS_{q-1})^{2^n}|CP_n\right]\left[(CS_{q-1})^{2^n}|LP_n\right]\left[(CS_{q-1})^{2^n}|RP_n\right]^{p-2} \end{aligned}$$

(b) Let n be even, thus the R-parity of CP_n is even as well and the R-parity of $\left((CS_{q-1})^{2^n}|CP_n\right)$ is also even. The first F-harmonic of $\left((CS_{q-1})^{2^n}|CP_n\right)$ is

$$H_F^{(1)}\left[(CS_{q-1})^{2^n}|CP_n\right] = \left[(CS_{q-1})^{2^n}|CP_n\right]\left[(CS_{q-1})^{2^n}|RP_n\right]$$

It is easy to see that $H_F^{(1)}\left[(CS_{q-1})^{2^n}|CP_n\right]$ has odd R-parity. As a consequence, on appending the expression $\left[(CS_{q-1})^{2^n}|CP_n\right]$ to $H_F^{(1)}\left[(CS_{q-1})^{2^n}|CP_n\right]$, in order to obtain the second F-harmonic, the C will be switched to L, keeping the same R-parity. The same will happen for the next F-harmonics, thus giving

$$\begin{aligned} H_F^{(p-1)}\left[(CS_{q-1})^{2^n}|CP_n\right] = \; & \left[(CS_{q-1})^{2^n}|CP_n\right]\left[(CS_{q-1})^{2^n}|RP_n\right] \\ & \left[(CS_{q-1})^{2^n}|LP_n\right]^{p-2} \end{aligned}$$

We have thus proven that, for even as well as for odd n, the same results are achieved both ways.

Let the R-parity of the q-periodic supercycle pattern be odd. In this case we will have to compute

$$\left((CS_{q-1})^{p\cdot 2^n}|\overline{CS_{p\cdot 2^n-1}}\right)$$

using Theorem 4. On the other hand, the n-th pitchfork bifurcation will be

$$\left((CS_{q-1})^{2^n}|\overline{CP_n}\right)$$

and it has the same R-parity as $\overline{CP_n}$ according to Lemma 2 in the Appendix. Furthermore $\overline{CP_n}$ has odd (even) R-parity when n is even (odd) as indicated in Note 1. Therefore, when F-harmonics are calculated, the same results will be obtained.◀

5.2 Convergence of first and last appearance orbits in an arbitrary window.

Theorem 6: *Inside any q-periodic window, the last apperance orbits within the $(n+1)$-chaotic band have the same accumulation point as the first appearance orbits within the n-chaotic band.*

Proof. Once again we study separately the cases of even and odd R-parity:

Let the R-parity of the q-periodic supercycle pattern be even. We want to compute the pattern of the supercycles of the $p\cdot 2^n$ first appearance and $p\cdot 2^{n+1}$ last appearance saddle-node orbits within the q-window. These orbits are located in the n- and $(n+1)$-chaotic bands of that window, respectively. In order to do so, we will apply Theorem 4, and hence we need to know the patterns of the $p\cdot 2^n$ first appearance and $p\cdot 2^{n+1}$ last appearance saddle-node orbits within the canonical window, which are located in the n- and $(n+1)$-chaotic bands, respectively.

(i) Let n be even.

The pattern of the supercycle associated to the first appearance $p\cdot 2^n$ saddle-node orbit, in the canonical window, is (see 10)

$$CS_{p\cdot 2^n-1} = CP_nRP_nLP_n(RP_n)^{p-3} \tag{13}$$

where $p = 3, 5, 7, \ldots$. Also, the pattern of the supercycle of the last appearance $p\cdot 2^{n+1}$ saddle-node orbit, in the canonical window, is given by (see 11)

$$CS_{p\cdot 2^{n+1}-1} = CP_nRP_nLP_n(RP_n)^{2\cdot p-3} \tag{14}$$

- The pattern of the supercycle associated to the $p \cdot 2^n$ first appearance saddle-node orbit, within the q-window, is computed by applying Theorem 4 as follows:
  (a) Repeat $p \cdot 2^n$ times the pattern CS_{q-1}
  $$(CS_{q-1})^{p \cdot 2^n}$$
  (b) Conjugate it according to (13)
  $$\left[(CS_{q-1})^{2^n}|CP_n\right]\left[(CS_{q-1})^{2^n}|RP_n\right]\left[(CS_{q-1})^{2^n}|LP_n\right] \left[(CS_{q-1})^{2^n}|RP_n\right]^{p-3} \tag{15}$$

- The pattern of the supercycle associated to the $p \cdot 2^{n+1}$ last appearance saddle-node orbit, within the q-window, by applying Theorem 4 is computed as:
  (a) Repeat $p \cdot 2^{n+1}$ times the pattern CS_{q-1}
  $$(CS_{q-1})^{p \cdot 2^{n+1}}$$
  (b) Conjugate it according to (14)
  $$\left[(CS_{q-1})^{2^n}|CP_n\right]\left[(CS_{q-1})^{2^n}|RP_n\right]\left[(CS_{q-1})^{2^n}|LP_n\right] \left[(CS_{q-1})^{2^n}|RP_n\right]^{2p-3} \tag{16}$$

For $p \to \infty$, the patterns (15) and (16) converge to the same Misiurewicz point with preperiod
$$\left[(CS_{q-1})^{2^n}|CP_n\right]\left[(CS_{q-1})^{2^n}|RP_n\right]\left[(CS_{q-1})^{2^n}|LP_n\right] \tag{17}$$
and period
$$\left[(CS_{q-1})^{2^n}|RP_n\right] \tag{18}$$

(ii) Let n be odd.

The pattern of the supercycle associated to the first appearance $p \cdot 2^n$ saddle-node orbit in the canonical window is (see 10)
$$CS_{p \cdot 2^n - 1} = CP_n LP_n RP_n (LP_n)^{p-3} \tag{19}$$
where $p = 3, 5, 7, \ldots$. Also, the pattern of the supercycle associated to the last appearance $p \cdot 2^{n+1}$ saddle-node orbit in the canonical window is given by (see 11)
$$CS_{p \cdot 2^{n+1} - 1} = CP_n LP_n RP_n (LP_n)^{2p-3} \tag{20}$$

- Pattern of the supercycle associated to the $p \cdot 2^n$ first appearance saddle-node orbit, within the q-window, by applying Theorem 4 is computed as:

  After repeating $p \cdot 2^n$ times the pattern CS_{q-1} and conjugating it according to (19) we obtain

  $$\left[(CS_{q-1})^{2^n}|CP_n\right]\left[(CS_{q-1})^{2^n}|LP_n\right]\left[(CS_{q-1})^{2^n}|RP_n\right] \left[(CS_{q-1})^{2^n}|LP_n\right]^{p-3} \tag{21}$$

- Pattern of the supercycle associated to the $p \cdot 2^{n+1}$ last appearance saddle-node orbit, within the q-window, by applying Theorem 4 is computed as:

  After repeating $p \cdot 2^{n+1}$ times the pattern CS_{q-1} and conjugating it according to (20) we obtain

  $$\left[(CS_{q-1})^{2^n}|CP_n\right]\left[(CS_{q-1})^{2^n}|LP_n\right]\left[(CS_{q-1})^{2^n}|RP_n\right] \left[(CS_{q-1})^{2^n}|LP_n\right]^{2p-3} \tag{22}$$

For $p \to \infty$, the patterns (21) and (22) converge to the same Misiurewicz point with preperiod

$$\left[(CS_{q-1})^{2^n}|CP_n\right]\left[(CS_{q-1})^{2^n}|LP_n\right]\left[(CS_{q-1})^{2^n}|RP_n\right] \tag{23}$$

and period

$$\left[(CS_{q-1})^{2^n}|LP_n\right] \tag{24}$$

◀

Let the R-parity of the q-periodic supercycle pattern be odd. Given that the R-parity of CS_{q-1} is odd, the conjugated patterns $\overline{CS_{p \cdot 2^n - 1}}$ and $\overline{CS_{p \cdot 2^{n+1} - 1}}$ will be used on applying Theorem 4. Hence, the results obtained will be the same.

Corollary to Theorem 3 is now rewritten, valid for any periodic window in the bifurcation diagram.

Corollary to Theorem 6. *Band merging points in a q-periodic window, whose patterns are given in (17)-(18) and (23)-(24) as the limit orbits of saddle-node orbit sequences, form a Misiurewicz point cascade. Furthermore, this Misiurewicz point cascade converges to the Myrberg-Feigenbaum point of that window, driven by the behavior of the saddle-node bifurcation cascades, described in [4].*

It is important to notice that the symbolic sequences of the Misiurewicz, given by (17-18) and (23-24), show the band merging points in any window, that is, not only those (already known) corresponding to the canonic window, but also those located inside every subwindow.

6 Discussion and conclusions

It has been shown, in the preceding sections, how the patterns of those periodic orbits of the Feigenbaum cascade determine the patterns of the orbits of the saddle-node bifurcation cascades. The accumulation points corresponding to these latter patterns are the band merging points, that are also Misiurewicz points. The sequence of these Misiurewicz points converges to the Myrberg-Feigenbaum point, and determines the chaotic band structure of the bifurcation diagram. This chain of relations shows how the periodic part of a window determines its chaotic part: the hyperbolic components of the set, determine the non-hyperbolic ones.

The relationship between the Feigenbaum cascade and the saddle-node bifurcation cascade could be expected, because both cascades share the same scaling law found by Feigenbaum [1]. On the other hand, this scaling is the direct consequence of the renormalization equation. This equation has only one fixed point in the functional space, whose stable manifold is one-dimensional. Thus, Feigenbaum's cascade and saddle-node cascade can but pertain to the same scaling universality [20].

The problem of localizing the chaotic band merging points, as well as the convergence of that sequence of points, is a problem largely addressed numerically by many authors as, for instance, by Hao and Zhang [21] in their "Bruselator oscillator" model. Both problems can be approached using the Corollary to Theorem 6, stated above, and the saddle-node cascade convergency, described in [4].

From the symbolic sequences of a pitchfork orbit in the Feigenbaum's cascade, the symbolic sequences of the supercycles of the saddle-node orbits can be computed. After which, Myrberg's formula [17] can be used to compute the value of the parameter giving rise to those orbits. This circumstance can be very useful to the experimenter, because once the Feigenbaum cascade has been observed, the corresponding saddle-node bifurcations can be spotted from it. Given that many dynamical systems, in particular lasers, present period doubling cascades, it easy for the experimenter to explicitly localize the saddle-node bifurcation cascades from the experimentally obtained return-map [22] . As a result, it is also possible to spot the intermittency cascades associated with that saddle-node bifurcation cascade. We must take into account that without an exact knowledge of the parameter values, it is practically impossible to localize these cascades, because of the geometrical stretching of those regions under saddle-node bifurcations.

We want to point out other possible uses from an experimental point of view. A relevant fact for the experimenter is the possibility to assess the localization of the Misiurewicz points which separate chaotic bands within any window. In this way, parameter values can be bound to the particular region of interest where the sought phenomena take place.

There is yet another interesting point. In all the results just obtained, orbit sequences have been considered. Each of them had a symbolic sequence associated to it. These symbolic sequences on their own, can be used to assess the underlying universal scaling laws [23]. This can be reflected, for instance, in the fact that the previously signaled Misiurewicz points scale according to the same Feigenbaum scaling scheme.

All that has been shown in this paper is immediately extendible to unimodal maps with extrema of the form x^{2n} under very general considerations.

Appendix

Lema 1 *Let $f(x;\, \mu)$ be an unimodal C^2 class map of the interval $[a,\, b]$ into itself, such that its critical point $x_c \in (a,\, b)$ is a maximum (minimum). If for $\mu = \mu_0$, there exists a q-periodic supercycle having a pattern CS_{q-1}, then $f^q(x_c;\, \mu_0)$ has a maximum (minimum) when the R-parity of CS_{q-1} is even, while it has a minimum (maximum) in case of odd R-parity of CS_{q-1}.*

Proof: Let us consider a value of μ in a small neighborhood of μ_0, such that $f^q(x;\, \mu)$ has a fixed point at some $x = x_c + \varepsilon$, for some arbitrarily small $\varepsilon > 0$. At $x = x_c$, $f^q(x_c;\, \mu)$ has a critical point, given that $f(x_c;\, \mu)$ has a critical point, for $x = x_c$. To determine whether it is a maximum or a minimum, it is enough to see whether, in a neighborhood of $x_c + \varepsilon$, the iterated function $f^q(x_c + \varepsilon;\, \mu)$ is increasing (corresponding to f having a minimum at x_c) or decreasing (corresponding to a maximum). That is, whether $[f^q]'\,(x_c + \varepsilon;\, \mu)$ is positive or negative.

$$[f^q]'\,(x_c + \varepsilon;\, \mu) = \prod_{i=1}^{q} f'(x_i)$$

where the x_i represent each of the q iterates of the function f period, starting at $x_1 = x_c + \varepsilon$. It turns out that

$$\operatorname{sign}\,[f^q]'\,(x_c + \varepsilon;\, \mu) = \prod_{i=1}^{q} \operatorname{sign}\, f'(x_i;\, \mu)$$

Let us assume f to have a maximum at x_c, that is, to be an increasing function in $(a,\, x_c)$ (i.e. with positive derivative) and a decreasing function in $(x_c,\, b)$ (i.e. with negative derivative). When f^q has a minimum at x_c (i.e. $\operatorname{sign}\,[f^q]'\,(x_c + \varepsilon;\, \mu)$ is positive) the number of x_i laying in $(x_c,\, b)$ will be even, therefore, the number of R-symbols in the pattern of the orbit will be even. Conversely, when f^q has a maximum at x_c, the number of R-symbols in the pattern will be odd.

Considering $x_1 = x_c + \varepsilon \to x_c$, we get back the superstable orbit, CS_{q-1}. This represents the deletion of the first R of the pattern, corresponding to x_1. Based on the continuous variation of the function with μ, we find that for an even R-parity of the supercycle pattern, a maximum will be present at $x = x_c$, while for an odd R-parity, there will be a minimum. ◀

Note 1: We need to know the R-parity of CP_n in order to compute F-harmonics of the pitchfork orbits in the proof. It is clear that for the period-1 orbit supercycle, whose sequence is C, its first pitchfork has the sequence CR and odd R-parity; the orbit originated by the second pitchfork has sequence $CR[C]R \to CRLR$, as given by the Metropolis-Stein-Stein harmonic [11], having even R-parity; the iteration of these harmonics gives odd R-parities for those CP_n with n odd, while those CP_n with even n, have even R-parity. As a consequence of this, it is easy to see that $\overline{CP_n}$ has odd (even) R-parity whenever n is even (odd).

Lemma 2 *The n-th pitchfork bifurcation of CS_{q-1}, that is, $CP_{n,q}$ is given by*

$$\left((CS_{q-1})^{2^n} | CP_n\right) \text{ for even } R\text{-parity of } CS_{q-1}$$

$$\left((CS_{q-1})^{2^n} | \overline{CP_n}\right) \text{ for odd } R\text{-parity of } CS_{q-1}$$

Furthermore, $CP_{n,q}$ has the same R-parity as CP_n (respectively, $\overline{CP_n}$) for even (respectively, odd) R-parity of CS_{q-1}.

Proof.

Let the R-parity of CS_{q-1} be even. When the MSS composition rule [11] is applied, the sequence S_{q-1} does not play any role, because it does not change the R-parity of the new pattern. Only the C's formerly changed have a role. Therefore we work as if with the n-th pitchfork bifurcation of the period-1 orbit.

Let the R-parity of CS_{q-1} be odd. The proof is similar to the former point, but the sequence S_{q-1} changes the R-parity of the new pattern when the composition rule is applied. Therefore we work as if with the conjugated of the n-th pitchfork bifurcation of the period-1 orbit.

Whatever the R-parity of S_{q-1} is, the $(CS_{q-1})^{2^n}$ sequence always has an even number of S_{q-1} sequences, therefore this even number of S_{q-1} gives an even R-parity. So, the R-parity of $\left((CS_{q-1})^{2^n}|CP_n\right)$ comes from the C's inserted from CP_n, that is, the R-parity of $\left((CS_{q-1})^{2^n}|CP_n\right)$ is that of CP_n. In the same way it is proven that $\left((CS_{q-1})^{2^n}|\overline{CP_n}\right)$ has the same R-parity as $\overline{CP_n}$.

References

[1] M. J. Feigenbaum. Quantitative universal for a class of nonlinear transformations. *J. Statist. Phys.*, 19:25–, 1978.

[2] M. J. Feigenbaum. The universal metric properties of nonlinear transformations. *J. Statist. Phys.*, 21:669–706, 1979.

[3] J. Milnor and W. Thurston. On iterated maps of the interval. In *Lect. Notes Math.*, number 1342, pages 465–563. Springer-Verlag, Berlin, New York, 1988.

[4] J. San Martín. Intermittency cascade. *Chaos, Solitons and Fractals*, 32:816–831, 2007.

[5] M. Misiurewicz and Z. Nitecki. Combinatorial patterns for maps of the interval. *Mem. Am. Math. Soc.*, 94:456, 1991.

[6] A. D. Churukian and D. R. Snider. Finding the windows of regular motion within the chaos of ordinary differential equations. *Phys. Rev. E*, 53:74–79, 1996.

[7] D. Xu and S. R. Bishop. Switching between orbits in a periodic window. *Phys. Rev. E*, 54:6940–6943, 1996.

[8] S. R. Bishop, D. Xu, C-Y Liau, and E-S Chan. Applying control in periodic windows. *Chaos, Solitons and Fractals*, 9:1297–1305, 1998.

[9] J. A. Blackburn and N. Gronbech-Jensen. Phase diffusion in a chaotic pendulum. *Phys. Rev. E*, 53:3068–3072, 1996.

[10] B. Derrida, A. Gervois, and Y. Pomeau. Universal metric properties of bifurcations of endomorphisms. *J. Phys A*, 12:269–296, 1979.

[11] N. Metropolis, M. L. Stein, and P. R. Stein. On finite limit sets for transformations on the unit interval. *J. Comb. Theory A*, 15:25–44, 1973.

[12] G. Pastor, M. Romera, and F. Montoya. On the calculation of Misiurewicz patterns in one-dimensional quadratic maps. *Physica A*, 232:536–553, 1996.

[13] M. Romera, G. Pastor, and F. Montoya. On the cusp and the tip of a midget in the Mandelbrot set antenna. *Phys Lett A*, 221:158–162, 1996.

[14] G Pastor, M. Romera, G. Alvarez, and Montoya F. Operating with external arguments in the Mandelbrot set antenna. *Physica D*, 171:52–71, 2002.

[15] G. Pastor, M. Romera, G. Alvarez, and F. Montoya. How to work with one-dimensional quadratic maps. *Chaos, Solitons and Fractals*, 18:899–915, 2003.

[16] T. Post and H.W. Capel. Windows in one-dimensional maps. *Phys. A*, 178:62–100, 1991.

[17] P. J. Myrberg. Iteration der reellen polynome zweiten grades III. *Ann. Acad. Sci. Fenn., Series A*, 336:1–18, 1963.

[18] W.-Z. Zheng, B.-L. Hao, G.-R. Wang, and S.-G. Chen. Scaling property of period n-tupling sequences in one-dimensional mappings. *Commun. Theor. Phys.*, 3:283–295, 1984.

[19] B. R. Hunt, J. A. C. Gallas, C. Grebogi, J. A. Yorke, and H. Kocak. Bifurcation rigidity. *Physica D*, 129:35–56, 1999.

[20] W. de Melo and S. van Strein. *One-dimensional dynamics.* Springer-Verlag, Berlin, 1993.

[21] B.-L. Hao and S.-Y. Zhang. Hierarchy of chaotic bands. *J. Stat. Phys.*, 28:769–, 1982.

[22] A. N. Pisarchik, R. Meucci, and F. T. Arecchi. Discrete homoclinic orbits in a laser with feedback. *Phys. Rev. E*, 62:8823–8825, 2000.

[23] R. López-Ruiz. Order in binary sequences and the routes to chaos. *Chaos, Solitons and Fractals*, 27:1316–1320, 2006.